

LIST OF FISHES COLLECTED AND OBSERVED IN NORTH  
AND SOUTH CAROLINA BY THE STEAMER "TIGER HAWK",  
DURING THE WINTER OF 1890-'91.

W. G. Woodruff, Naturalist.

*Percopygion concolor.*

One specimen from Winyah Bay. Taken in  
June 1891.

*Micropterus salmoides.*

Many were taken by the fishermen in large  
schools at Cape Lookout, N. C.

*Sciaenops ocellatus.*

Very common with *M. salmoides* at Cape Look-  
out. Also caught occasionally on  
head lands at other places in rivers.

Very abundant at Cape Lookout.

*Dasistis sayi.*

One specimen taken at Hattaway Bay Is. 2.

Edisto R., many at Cape Lookout, N.C.

*Acipenser sturio* var. *oxyrhynchus* ?

Abundant in rivers. Fishermen take many  
in drift nets. Caviar is made from  
the roe.

*Lepidosteus osseus.*

Very plentiful in fresh rivers. So  
abundant that shad fishing is al-  
most impossible as the bar is ruin-  
ous to the nets. Taken in South  
Edisto R. and Winyah Bay.

*Acia calva.*

Abundant in the vicinity of Georgetown,  
S. C.

*Amincus platycerhalus.*

Taken in Sampit river at Georgetown.  
Stomach full of filth.

*Hybernathus nuchalis.*

Taken in numbers in the Sampit R. at  
Georgetown, S. C., December 30, 1892.

*Clupea sapidissima.*

Taken in the South Edisto River and Win-  
yah Bay, February 28th and March 31,  
respectively.

*Opiathonus chinns ?*

Young found plentifully at Chisholm's Id.,  
Crossw River.

*Clupea medius*

South Edisto R. and Winyah Bay with *C.*  
*sapidissima.*

*Brevortia tyrannus.*

South Edisto R. February 28. Winyah Bay  
March 31. In both localities the  
fish were affected with the parasite  
*Cymothoa praeputator* and a *fenax*.

~~*Stolephorus Mitchilli.*~~

*Stalophorus mitchilli.*

Tybee Roads, Ga., March 13, 1891; Vero-  
beach City, N. C., February 21, 1891.  
Not very abundant.

*Pandulus nebulosus.*

North Island, Winyah Bay, January 13, '91;  
Tybee Roads, Ga., March 13, 1891;  
Skull Creek, S. C.

*Pandulus heteroclitus.*

North Island, Winyah Bay, January 6, 1891;  
plentiful. Tybee Roads, Ga., March  
15, 1891; Skull Creek, Edisto Island,  
March 3, 1891; Ashley River, Coosaw  
River, February 19. This seems to  
be the most abundant species of Cy-  
prinodont in this region.

*Cyprinodon variegatus.*

N. Island, Winyah Bay, January 6, 1891.  
A few small ones taken.

*Mucenia parva.*

North Island, Winyah Bay, January 6, 1891;

plentiful at this place.

*Melipotis latipinna.*

A few found at North Island, Winyah Bay

and at Edisto Island, March 3, 1891.

*Gambusia patricellia.*

Lady's Island, North Island, Edisto Island

*Rafes reticulata.*

Said to be abundant in the Pee Dee River.

Rare a few at this place.

*Anguilla rostrata.*

Small ones taken at Georgetown, December

20, 1890 and at Georgetown Light,

Winyah Bay.

*Siphonura louisianae.*

Winyah Bay, Feb. 29, 1891. Morehead

City, N. C., April 7, 1891.

*Migil cephalus.*

Winnah Bay and Turrit River, very abundant. Few taken December 20, 1890. Many about four inches long.

*Quarimus giranis.*

Very abundant in Winnah Bay in January '91. Few taken all along the coast later in the season.

*Menidia notata.*

Abundant all along the coast all winter. Spawning females taken April 7, 1891, at Morehead City, N. C. No males observed here.

*Menidia laciniata.*

Taken April 7, 1891 at Morehead City, N. C. This species seems to be all males, associated with *M. notata* which so far as was observed were all females.

*Menidia peninsulae.*

Taken at Georgetown, December 20, 1890.

*Trichinurus lepturus.*

One taken at Katteras Inlet.

*Pogonocherus saltatrix.*

Very many small ones taken in larval salines  
at Cape Lookout, E. C., April 6, '91.

*Stromateus triacanthus.*

Katteras Inlet, N. C. Fort Mason, N. C.

*Stenobrycon pulchus.*

Pee Dee River, December 31, 1890.

*Lepomis pulchus.*

Abundant in Pee Dee River. Taken Dec.  
31, 1890.

*Lepomis gibbosus.*

Pee Dee River, December 31, 1890.

*Micropterus salmoides.*

Abundant Pee Dee River.

*Morone americana.*

Pee Dee River, Dec. 31, 1890, Sampit R.  
December 20, 1890.

*Murexana atrarius.*

Wales Branch, S. O.

*Locostes rhomboides.*

very plentiful at Incehead City, Apr. 7, '91.

*Pezomachus chionis.*

Young taken at Tybee Roads, very plentiful.

*Psiridialis chryseus.*

Tybee Roads- abundant. Incehead Quay,  
Feb, April 7, 1891.

*Teucha caelata.*

Abundant all along the coast.

*Mioscirtus xanthopus.*

Abundant all along the coast. Young  
taken at Incehead City, April 7, '91.

*Psiridialis alternans.*

Patience Inlet. Type lost.



*Cynoscopus regalis.*

Common at Cape Lockout. Common there  
in large numbers in large quantities.  
This species here seems to take the  
place of the *C. maculatus* of farther  
south, although both species are  
found all along the coast.

*Astrascopus anoplus.*

Whisholas Island, Coosaw River, Feb. 15, '91  
North Island, Winyah Bay, Jan. 9, '91  
Coosaw River February 12, '91. Not  
uncommon. All years.

*Habius paleosoma.*

Saville River, Dec. 30, 1890; Winyah Bay,  
December 31, 1890.

*Coliasa hysci.*

Charleston Harbor, Skull Creek. May 11.  
Jericho Creek. Fort Royal Island.  
Parrot Creek. Abundant everywhere.

*Prionotus*

*Trinectes laticollis*.

Charleston Inlet, Cape Fear River, Marshwood  
City, abundant.

*Gobiosoma aeneum*.

Cooper R. North Island. Not uncommon.

*Parablennius latipes*.

Abundant everywhere. Young taken in  
quantities during whale season.

*Parablennius aeneus*.

Jericho Creek, Fort Royal Tavern, not  
at all rare.

*Parablennius aeneus*.

Jericho Creek, Station 567, not all  
green, common.

*Parablennius aeneus*.

Not found on Cape Fear River Inlet.

*Parablennius aeneus*.

Tybee Island, young; North Island, young;  
Charleston Harbor, young.

*Paralichthys dentatus.*

Winyah Bay, Charleston Harbor.

*Paralichthys albigutta.*

Winyah Bay.

*Paralichthys lethostigma.*

Geos. W. River, Parrot Creek Y.

*Etroplus crenolatus.*

Sea saw River, February 18, 1891. Parrot  
Creek, Callabogue Sound, and other  
places. Not rare.

*Ancylorhiza quadracelata.*

Geos. W. River, February 18, 1891. Callabor-  
ne Sound, N. Inlet. Not very rare.

*Achirus fasciatus.*

Charleston Harbor. Abundant.

*Aphoristicia plagiosa.*

Abundant everywhere.

*Chloroceryle geometricus.*

Lowland City; Cape Lookout; very common.